Vol 4 Issue 4 May 2014

ISSN No: 2230-7850

## International Multidisciplinary Research Journal

# Indian Streams Research Journal

Executive Editor Ashok Yakkaldevi

Editor-in-Chief H.N.Jagtap

#### Welcome to ISRJ

#### RNI MAHMUL/2011/38595

ISSN No.2230-7850

Indian Streams Research Journal is a multidisciplinary research journal, published monthly in English, Hindi & Marathi Language. All research papers submitted to the journal will be double - blind peer reviewed referred by members of the editorial board. Readers will include investigator in universities, research institutes government and industry with research interest in the general subjects.

#### International Advisory Board

Flávio de São Pedro Filho Federal University of Rondonia, Brazil

Kamani Perera

Regional Center For Strategic Studies, Sri

Lanka

Janaki Sinnasamy

Librarian, University of Malaya

Romona Mihaila

Spiru Haret University, Romania

Delia Serbescu Spiru Haret University, Bucharest, Romania

Anurag Misra

DBS College, Kanpur

Titus PopPhD, Partium Christian University, Oradea, Romania

Mohammad Hailat

Dept. of Mathematical Sciences, University of South Carolina Aiken

Abdullah Sabbagh Engineering Studies, Sydney

Catalina Neculai University of Coventry, UK

Ecaterina Patrascu

Spiru Haret University, Bucharest

Loredana Bosca

Spiru Haret University, Romania

Fabricio Moraes de Almeida Federal University of Rondonia, Brazil

George - Calin SERITAN

Faculty of Philosophy and Socio-Political Sciences Al. I. Cuza University, Iasi

Hasan Baktir

English Language and Literature

Department, Kayseri

Ghayoor Abbas Chotana Dept of Chemistry, Lahore University of

Management Sciences[PK]

Anna Maria Constantinovici AL. I. Cuza University, Romania

Horia Patrascu Spiru Haret University, Bucharest,Romania

Ilie Pintea,

Spiru Haret University, Romania

Xiaohua Yang PhD, USA

.....More

#### Editorial Board

Pratap Vyamktrao Naikwade Iresh Swami

ASP College Devrukh, Ratnagiri, MS India Ex - VC. Solapur University, Solapur

Head Geology Department Solapur

University, Solapur

Rama Bhosale Prin. and Jt. Director Higher Education,

Panvel

R. R. Patil

Salve R. N.

Department of Sociology, Shivaji University, Kolhapur

Govind P. Shinde

Bharati Vidyapeeth School of Distance Education Center, Navi Mumbai

Chakane Sanjay Dnyaneshwar Arts, Science & Commerce College,

Indapur, Pune

Awadhesh Kumar Shirotriya Secretary, Play India Play, Meerut (U.P.)

N.S. Dhaygude Ex. Prin. Dayanand College, Solapur

Narendra Kadu

Jt. Director Higher Education, Pune

K. M. Bhandarkar

Praful Patel College of Education, Gondia

Sonal Singh

Vikram University, Ujjain

G. P. Patankar

Maj. S. Bakhtiar Choudhary Director, Hyderabad AP India.

S.Parvathi Devi

Ph.D.-University of Allahabad Sonal Singh,

Vikram University, Ujjain

Rajendra Shendge

Director, B.C.U.D. Solapur University,

Solapur

R. R. Yalikar

Director Managment Institute, Solapur

Umesh Rajderkar

Head Humanities & Social Science YCMOU, Nashik

S. R. Pandya

Head Education Dept. Mumbai University,

Mumbai

Alka Darshan Shrivastava

S. D. M. Degree College, Honavar, Karnataka Shaskiya Snatkottar Mahavidyalaya, Dhar

Rahul Shriram Sudke Devi Ahilya Vishwavidyalaya, Indore

S.KANNAN Annamalai University,TN

Satish Kumar Kalhotra

Maulana Azad National Urdu University

Address:-Ashok Yakkaldevi 258/34, Raviwar Peth, Solapur - 413 005 Maharashtra, India Cell: 9595 359 435, Ph No: 02172372010 Email: ayisrj@yahoo.in Website: www.isrj.net

Indian Streams Research Journal ISSN 2230-7850 Volume-4 | Issue-4 | May-2014 Available online at www.isrj.net





## FLORISTIC ENUME RATION OF SRI VENKATESWARA ZOOLOGICAL PARK FOR THE FUTURE CONSERVATION OF BIODIVERSITY

#### A. Job Roger Binny and N. Yasodamma

Dept of Botany, Sri Venkateswara University, Tirupati .

Abstract:-Sri Venkateswara Zoological Park is a protected area of an ex-situ conservation for certain fauna and in-situ conservation of natural floral biodiversity comes under Sri Venkateswara Biospere Reserve Forest situated in the Seshachalam Hill ranges of Southern Eastern Ghats of Peninsular India. Floristic studies of protected areas is very much essential to enumerate the plant species for future generations and to maintain Sustainable Biodiversity Conservation. The systematic florist studies for three years resulted a total of 802 plant species include Bryophytes (2), Pteridophytes (13), Gymnosperms (5) along with major taxa of angiosperms (782) includes dicotyledons 639 and monocotyledons 143. Distribution of species to that of the other districts and regional floras showed a great diversity in the ratio of dicots to monocots equally to that of Kurnool and Chittoor Districts floras. Species diversity is closely related with Anantapur, Kurnool, Chittoor districts as the largest genera Euphorbia (15), Justicia (12) and Cyperus (10) and also with largest family Leguminosae (108) (Fabaceae). It also represents red listed species in natural conditions with Pterocarpus santalinus, Santalum album along with 11 other threatened category species in the zoo park. Hence, it is very essential to enumerate and enlist the flora of a protected area for the sustainable conservation for future generation.

 $\textbf{Keywords:-} protected area, sustainable, biodiversity, Pterocarpus santalinus, Santalu \, album, threatened. \\$ 

#### INTRODUCTION

Protected areas are the key tools for the conservation of Biological diversity as established in the Convention on Biological Diversity. IUCN defines "the protected area as aclearly defined geographical space, recognized, dedicated and managed through legal or other effective means to achieve the long term conservation of nature with associated ecosystem services and cultural values". National parks are the protected areas come under Category II of CBD. Zoological parks are established as the centers for conservation, training and education with the main purpose of protection, study and research on wild life and education on biodiversity. Ex-situ conservation of biological diversity outside the natural habitats includes plant and animal collections, botanical gardens, zoological parks, wild life research facilities and germ plasm collections of wild and domestic taxa. A zoological park is principally of an environmental in nature committed to high quality conservation will be authorized to continue their activities, to understand the value of biodiversity of wild flora and fauna, ecosystems and the interdependence of all living organisms on earth including human being. Man's survival is intimately related with availability of the plant wealth and also acquiring scientific knowledge about the flora and vegetation of a region is essential with emphasis on distribution and status of rare and endangered species for future generations.

All protected areas should aim to conserve the composition, structure, function and evolutionary potential of Biodiversity, contribute to regional conservation strategies as core reserves, buffer zones, corridors, stepping stones for majority of species, to maintain diversity of land scape or habitat, associated species and ecosystems; geomorphology and geology; provide regulatory ecosystem services, including buffering against the impacts of climate change; conserve natural and scenic areas of national and international significance for cultural, spiritual and scientific purposes; facilitate low impact scientific research activities and ecological monitoring related to consistent with the values of the protected area; and help to provide educational opportunities in the management approaches (CNPPA & IUCN., 1992; 1993; 1994).

A. Job Roger Binny and N. Yasodamma , "FLORISTIC ENUME RATION OF SRI VENKATESWARA ZOOLOGICAL PARK FOR THE FUTURE CONSERVATION OF BIODIVERSITY" Indian Streams Research Journal | Volume 4 | Issue 4 | May 2014 | Online & Print

#### ABOUT THE SELECTED STUDY AREA-S.V. ZOOLOGICAL PARK: (Fig 1-3)

The study area Sri Venkateswara Zoological Park is a protected area for in-situ conservation of natural flora and exsitu conservation of certain fauna as a part of conservation of Biodiversity. Sri Venkateswara Zoological Park is situated between the latitudes of 130 37' 59.52" to 130 37' 33.6 " North and longitudes of 790 21' 7.28" to 790 21' 43.2" East with an average elevation of 749 feet MSL altitudinal range. S.V. Zoological park is surrounded by Tirupati towards North, Cherlopalli bypass road and Regional Science Centre towards East, Sreevari mettu on Western side and Tirumala, Kalyanidam, Cherlopalli and Pudipatla towards South as boundaries. Zoo park area generally comprises with daily temperatures ranging from 160 C – 450 C, and an average annual rainfall about 934 mm received mainly from North east monsoon. S.V Zoological Park is a part of S.V. Wild life sanctuary of S.V. National park in the Seshachalam Hill Ranges of Southern Eastern Ghats of Peninsular India with geographical area of 5532 acres (2212 Ha) (22 sq km). The Zoo Park has been established on the modern concepts of Zoo management of protected areas founded on 29.09.1989 and opened to the public on 30.04.1993 to facilitate captive breeding of the endangered and threatened animals and rehabilitate them with a special focus on endemic fauna and educate visitors about wildlife conservation through mythological theme nearer to the natural habitat. (http://forest.ap.nic.in/SVZP\_Introduction. htm).

The Zoo Park is developed into enclosures, enclaves and Moats named Kishkindha vanam Angada Dweep-Macaca speciosa (Stumptailed macaque), Vali Dweep-Macaca mulatta (Rhesus macaque), Sugreeva Dweep-Macaca radiata (Bonnet macaque), Karthikeya vanam or Mayur Vanam-Pavo cristatus(Common Peafowl), Anthropoides virgo(Demoiselle Crane), Swanra Harini Vanam or Mareecha Vanam-Axis axis (Chital or Spotted deer), Chitra Koota Vanam-Perdicula asiatica (Jungle bush quail), Galloperdix lunulata (Painted spur fowl), Francolinus pondicerianus (Grey Partridge), Lophura nycthemera (Silver pheasant), Suka Nivas or Vihanga Vanam or ChitraKoota Vanam-Gallus soneratti (Grey junglefowl), Psittacula krameri (Roseringed parakeet), Psittacula cyanocephala (Blossom Headed Parakeet), Pistlacula himalayana (Slaty Headed Parakeet), Placeus phillippinus (Indian Baya), Kanithi Vanam-Cervus unicolor (Sambar), Gokulam-Boselaphus tragocamelus (Nilgai or Blue bull), Krishna Mruga Vanam-Antilope cervicapra (The Blackbuck or The Indian antelope), Mruga Vanam-Tetracerus quadricornis (The Fourhorned Antelope), Grasa Mruga Vanam-Gazella gazella (Chinkara), Vruka Vanam-Canis lupus linnaeus (Wolf), Bhagela Vanam-Panthera pardus (Leopard), Shakti Vanam-Panthera tigris (Linnaeus) (The tiger), Bharat Vanam-Panthera leo (Linnaeus) (The Lion), Vyaghra Vanam-Panthera tigris tigris (White Tiger), Dandaka aranya-Hyaena hyaena (Striped Hyena), Damanaka Vanam-Vulpes bengalensis (Indian fox), Shunaka Vanam-Cuon alpinus (Dhole), Karataka Vanam-Canis aureus linnaeus (Jackal), Hystrix indica (The Indian Porcupine), Jambavantha Vanam-Melursus ursinus ( Sloth Bear), Varaha Vanam-Sus scrofa (Wild Boar), Nala Neela Dweepam-Theropithecus gelada (Baboon), Airavatha Vanam-Elephas maximus (The Indian Elephant), Makara Nivas-Crocodylus palustris (Marsh crocodile), Emu-Dromaius novaehollandiae (Emu), Neela Vanam-Common Langur, Koorma Nivas-Gochelone elegans (Starred tortoise). Melanochelys trijuga (Indian pond terrapin), Manasa Sarovar-Ardea purpurea (Purple Heron), Mycteria leucocephala (Painted Stork), Threskiornis aethiopica (White Ibis), Anser indicus (Barheaded Goose), Tadorna ferruginea (Brahminy Duck), Anas acuta (Pintail), Anas creca (Common Teal), Nettapus coromandeliannus (Cotton Teal), Dendrocygna javanica (Lesser Whistling Teal), Sarkidiornis melanotos (Comb Duck), Amaurornis phoenicurus (White Breasted Water Hen), Porphyrio (Purple Moorhen), Anser owser (Greylag Goose), Anas querquedula (Garganey), Anasclypeata (Shoveller), Kalindi Sarovar-Pelecanus phileppensis (roseus), (Spotted billed pelican or Grey pelican), Padma Kheli Sarovar-Ciconia episcopus (White-necked stork).




Figure: I: S.V. Zoological Park (Founded 1989; open to public 1993)

Latitudes 130 371 59.5211 to 130 371 33.6 11 North Longitudes 790 211 17.2811 to 790 211 43.2 11 East Altitudinal range: 749 feet MSL; Area: 553 acres (22 sq.km).



Fig-2: Zoological Park located in S.V.National Park of Southern Eastern Ghats



#### Fig-3: Boundaries of Sri Venkateswara Zoological Park

#### MATERIALS AND METHODS:

#### Field work, herbarium preparation and Identification of Taxa:

Intensive field studies were undertaken during different seasons of every year from 2010 to 2013 for a period of 3 years in the S.V. Zoological park, collected the specimens systematically and prepared the herbarium as per the standard method (Jain and Rao (1977) and identified the species following Bentham and Hooker System of Classification with the help of the Flora of presidency of Madras, Flora of Tamilnadu Carnatic, Flora of Nallamalai Hills, Flora of Andhra Pradesh, Flora of Eastern Ghats, Flora of Kurnool, Anantapur, Chittoor and Nellore Districts and also identifications were confirmed, after matching with authentic specimens of the Herbarium, Department of Botany (SVUTY), Sri Venkateswara University, Tirupati. (Gamble and Fisher 1915-1936; Matthew K.M, 1981-1988., Ellis 1966., Venkata Raju and Pullaiah 1995., Pullaiah and Yesoda 1989., Rangacharyulu 1991., Madhava Chetty et al 2008., Suryanarayana and Srinivasa Rao., 2002).

#### **Nomenclature:**

It was tallied with the latest accepted names, author citations important synonyms and families with "The Plant List Version 1, 2010". The representative voucher specimens were numbered and deposited in the herbarium (SVUTY) Department of Botany S. V. University Tirupati.

#### Floristic Analysis:

A critical analysis has been made with the floristic studies reported from Seshachalam hill ranges. All the taxa were tabulated under four plant groups: Dicotyledons, Monocotyledons, Pteridophytes and Gymnosperms. Families with number of Genera and Species, the Dominant Families with Genera and Species were compared with other regional and district floras. Herbarium specimens 2615 voucher numbers were deposited in the Department of Botany, S. V. University, Tirupati.

#### **RESULTS:** (Table: 1-8) (Graphs: 1-3):

Floristic enumeration of the selected area resulted 802 species include Bryophytes (2), Pteridophytes (13), and Gymnosperms (5) along with the Angiosperms (782) with Dicotyledons (639) and Monocotyledons (143). Angiosperms species represented a total of 108 families; Monocots (20), Dicots (88) with monocotyledons to dicotyledon species ratio 1:4.47; genera ratio 1:4.66, family ratio 1:4.40 designated its uniformity in the diversity of monocotyledon species, genera and family. The vegetation type of Zoo denotes the dry deciduous spiny scrub forest dominates with grasses with spiny shrubs and tree species in its overall distribution, Major family represents Leguminosae (108 species Fabaceae) equally to that of the Madras Presidency and flora of Andhra Pradesh. Next largest family Poaceae 68 species is also closely related with the floras of Chittoor, Anantapur, Kurnool and Nellore districts. Top 10 genera more than 10 species; as the largest genus Euphorbia with 15 species flowed by Justicia 12, Cyperus 10 to that of Anantapur, chittoor and Kurnool districts. S.V. Zoological Park also comprises IUCN Red Listed species of Endangered Category 3, vulnerable 7, near Threatened 1, least concerned 94 and not evaluated with one species.

Table: 1: Floristic Distribution of Genera and Species of Sri Venkateswara Zoological Park Bentham & Hooker System of Classification: (1862- 1883)

		ICOT	ΓYΙ	LEDONS					GYMNOSPERMS	5		
	POL YPETALAE				S.No	Family	G	S	S.No	Family	G	S
S.N	Family	G	S		49	Passifloraceae	1	1	1	Araucariaceae	1	1
1	Magnoliaceae	1	1		50	Caricacaea	1	1	2	Cupressaceae	1	1
2	Annonaceae	3	4		51	Cucurbitaceae	12	14	3	Cycadaceae	1	3
3	Menispermaceae	4	4		52	Cactaceae	2	2		Total	3	5
4	Nymphaeaceae	1	2		53 Aizoaceae		4	6				
5	<ul> <li>a) Nelumbonaceae</li> </ul>	1	1		54 Apiaceae		1	1		MONOCOTYLEDO	NS	
6	Papaveraceae	1	1		55	Araliaceae	2	2	S.No.	Family	G	S
7	Capparidaceae	4	6	l		Total	176	285	1	Orchidaceae	2	2
8	Violaceae	1	1						2	b) Marantaceae	1	1
9	Bixaceae	1	1			GAMOPETALA			3	c) Cannaceae	1	1
10	Cleomaceae	1	5	l	S.No.	Family	G	S	4	d) Musaceae	1	1
11	Polygalaceae	1	5	ı	1	Rubiaceae	21	31	5	Amaryllidaceae	2	2
12	Caryophyllaceae	1	2		2	Asteraceae	36	36	6	Dioscoreaceae	1	2
13	Portulacaceae	2	5	ı	3	Plumbaginaceae	1	1	7	Roxburgiaceae	1	1
14	Clusiaceae	1	1		4	Sapotaceae	2	2	8	Liliaceae	2	2
15	Malvaceae	9	15		5	Ebenaceae	1	5	9	a) Asparagaceae	7	12
16	a) Bombacaceae	1	1	ı	6	Oleaceae	4	7	10	b) Colchicaceae	1	1
17	Sterculiaceae	6	6	l	7	Salvadoraceae	1	1	11	c) Hypoxidaceae	1	1
18	Tiliaceae	3	13	ı	8	Apocynaceae	12	15	12	Xyridaceae	1	1
19	Elaeocarpaceae	1	1		9	Asclepiadaceae	17	22	13	Commelinaceae	3	8
20	Linaceae	1	1		10	Loganiaceae	1	2	14	Arecaceae	7	7
21	a) Erythroxylaceae	1	1		11	Gentianaceae	1	1	15	Typhaceae	1	1
22	Malpighiaceae	2	2		12	Hydrophyllaceae	1	1	16	Araceae	1	1
23	Zygophyllaceae	3	3	l	13	Boraginaceae	6	10	17	Aponogetonaceae	1	2
24	Geraniaceae	1	1		14	Convolvulaceae	8	17	18	Eriocaulaceae	1	2
25	a) Oxalidaceae	2	2		15	Solanaceae	5	9	19	Cyperaceae	9	27
26	Rutaceae	8	9	4	16	Scrophulariaceae	4	7	20	Poaceae	40	68
27	Simaroubaceae	2	2	1	17	Bignoniaceae	8	12		Total	84	143
28	Ochnaceae	1	1	4	18	Pedaliaceae	3	6				
29	Burseraceae	2	3	ł	19	Acanthaceae	20	39				
30	Meliaceae	2	2	4	20	Verbenaceae	10	14				
31	Celastraceae	3	4	ł	21	Lamiaceae	11	21		Distribution of		
32	Rhamnaceae	3	5	ł		Total	173	259	В	ryophytes & Pteridop	hytes	
33	Vitaceae	3	4	ł		140110 OTT 114TD						
	Sapindaceae	4	5	1	C M-	MONOCHLAMYD			C M-	BRYOPHYTES		- C
35 36	Anacardiaceae Moringceae	6	6	ł	S.No.	Family	<b>G</b>	<b>S</b>	S.No.	Family Ricciaceae	G	<b>S</b>
37		27	59	1	2	Nyctaginaceae	11	19	2		1	1
38	a) Fabaceae b) Caesalpiniaceae	12	26	ł	3	Amaranthaceae Aristolochiaceae	11	19		Funariaceae <b>Total</b>	2	2
39	c) Mimosaceae	10	23	1	4	Lauraceae	1	1		1 Otai		
40		10	23	1	5					DTE DIDODUATE	C	
40	Rosaceae Droseraceae	1	1	1	6	Hernandiaceae Proteaceae	1	2	S.No.	PTERIDOPHYTE:	G	S
41	Combretaceae	3	8	1	7	Loranthaceae	2	2	5.No.	Dryopteridaceae	1	1
43	Myrtaceae	3	4	1	8	Santalaceae	2	2	2	Marsileaceae	1	1
44	Melastomaceae	1	1	$\vdash$	9	Euphorbiaceae	22	52	3	Nephrolepidaceae	1	3
45	Lythraceae	3	3	$\vdash$	10	a) Ulmaceae	22	2	4	Polypodiaceae	1	1
45	Onagraceae	1	1	$\vdash$	11	c) Moraceae	2	6	5	Polypodiaceae Pteridaceae	4	5
47	Flacourtiaceae	2	2	$\vdash$	12	Casuarinaceae	1	1	6	Salviniaceae	2	2
48		1	1	$\vdash$	12				0			
48	Turneraceae	1	1		L	Total	49	95		Total	10	13

Table: 2: Floristic enumeration of S.V. Zoological Park

S.No.	Division	Family	Genera	Species		
1	Bryophytes	2	2	2		
2	Pteridophytes	6	10	13		
3	Gymnosperms	3	3	5		
		•		•		
Al	NGIOSPERMS	108	482	782		
4	Dicotyledons	88	398	639		
5	Monocotyledons	20	84	143		
	Total	119	497	802		

Graph: 1: Floristic enumeration of S.V. Zoological Park




Table: 3. Percentage among Angiosperm families, genera and species

Taxa	Species	%	Genus	%	Family	%
Dicotyledons	639	81.71	398	82.57	88	81.48
Monocotyledons	143	18.29	84	17.43	20	18.51
Total	78	32	4	82	10	08

Table: 4. Ratios between Monocotyledon vs Dicotyledons families, genera and species

Taxa	Family	Genus	Species							
Dicotyledons	88	398	639							
Ratio		1:4.52:7.26								
Monocotyledons	20	143								
Ratio		1: 4.2 : 7.15								
Total	108 482 78									
Dicot/ Monocot		1:4.40:4.74:4.47								

Table-5: Distribution of major Angiosperm families, genera and species

	3. a. Top 20 Fa	milies			3.b.	Top 20 Genera	with more than 5 Sp	ecies	
S.No.	Family	Gen	Gen Sps S. No. Genus		Genus	Family	No. of Sps		
1	Leguminosae	49	108		1	Euphorbia	Euphorbiaceae	15	
2	Poaceae	40	68		2	Justicia	Acanthaceae	12	
3	Euphorbiaceae	22	52		3	Cyperus	Cyperaceae	10	
4	Acanthaceae	20	39		4	Acacia	Mimosaceae		
5	Asteraceae	36	36		5	Crotolaria	Fabaceae	9	
6	Rubiaceae	21	31		6	Senna	Caesalpiniaceae	9	
7	Cyperaceae	9	27		7	Fimbristylis	Cyperaceae		
8	Asclepiadaceae	17	22		8	Indigofera	Fabaceae	8	
9	Lamiaceae	11	21		9	Eragrostis	Poaceae		
10	Amaranthaceae	11	19		10	Ipomoea	Convolvulaceae	7	
11	Convolvulaceae	8	17		11	Phyllanthus	Euphorbiaceae	7	
12	Apocynaceae	12	15		12	Alysicarpus	Fabaceae		
13	Malvaceae	9	15		13	Barleria	Acanthaceae	6	
14	Cucurbitaceae	12	14		14	Corchorus	Tiliaceae	0	
15	Verbenaceae	10	14		15	Terminalia	Combretaceae		
16	Tiliaceae	3	13		16	Acalypha	Euphorbiaceae		
17	Bignoniaceae	8	12		17	Cleome	Cleomaceae		
18	Asparagaceae	7	12		18	Commelina	Commelinaceae	5	
19	Boraginaceae	6	10		19	Diospyros	Ebenaceae		
20	Rutaceae	8	9		20	Ficus	Moraceae		

Graph: 2: Distribution of Major Angiosperm Families, Genera and species




Table-6: Comparative Account of Dominant Families with Flora British India and Regional Floras

	6.a. Com	parisio	on of Top 10 Angiosp	erm F	amilies with regiona	ıl Floras	s of India				
S. No	S. V. Zoo Park	Sps	Madras Presidency Gamble (1915- 1935)	Sps	Flora of India (B.S.I., Jain, 1989)	Sps	Flora of A.P. T. Pullaiah et al.	Sps			
1	Leguminosae	108	Leguminosae	432	Poaceae	1225	Leguminosae	299			
2	Poaceae	68	Graminae (Poaceae)	388	Orchidaceae	990	Poaceae	281			
3	Euphorbiaceae	52	Rubiaceae	226	Leguminosae	775	Cyperaceae	139			
4	Acanthaceae	39	Acanthaceae	202	Asteraceae	754	Euphorbiaceae	126			
5	Asteraceae	31	Euphorbiaceae	199	Rubiaceae	495	Acanthaceae	108			
6	Rubiaceae	27	Orchidaceae	199	Cyperaceae	499	Asteraceae	105			
7	Cyperaceae	22	Compositae (Asteraceae)	189	Euphorbiaceae	419	Rubiaceae	88			
8	Asclepiadaceae	21	Cyperaceae	172	Lamiaceae	393	Orchidaceae	69			
9	Lamiaceae	21	Labiatae (Lamiaceae)	128	Acanthaceae	379	Lamiaceae	65			
10	Amaranthaceae	19	Asclepiadaceae	93	Schrophulariaceae	356	Convolvulaceae	61			
6.b. Comparision of Top 10 Dominant Families with Chittoor District Floras											
S.	S. V. Zoo	Sps	Ardhagiri		Chittoor		Chittoor				
No	Park	_	Hemalatha	Sps	Chetty et al.	Sps	D. Rangachari	Sps			
1	Leguminosae	108	Leguminosae	43	Leguminosae	215	Leguminosae	141			
2	Poaceae	68	Poaceae	20	Poaceae	150	Euphorbiaceae	58			
3	Euphorbiaceae	52	Euphorbiaceae	19	Asteraceae	82	Poaceae	53			
4	Acanthaceae	39	Malvaceae	14	Euphorbiaceae	80	Acanthaceae	42			
5	Asteraceae	31	Asteraceae	13	Acanthaceae	60	Cyperaceae	40			
6	Rubiaceae	27	Acanthaceae	13	Cyperaceae	58	Compositae (Asteraceae)	34			
7	Cyperaceae	22	Amaranthaceae	11	Rubiaceae	57	Rubiaceae	33			
8	Asclepiadaceae	21	Lamiaceae	11	Lamiaceae	54	Labiatae (Lamiaceae)	25			
9	Lamiaceae	21	Cyperaceae	11	Malvaceae	42	Malvaceae & Convolvulaceae	24			
10	Amaranthaceae	19	Rubiaceae	10	Convolvulaceae	37	Asclepiadaceae	20			
	6.c. Comparision	of To	p 10 Dominant Fam	ilies w	ith other Districts fl	oras of A	Andhra Pradesh				
S.	S. V. Zoo		Nellore		Anantapur	No.	Kurnool				
No	Park	Sps	Suryanarayana & Srinivasa Rao	Sps	T. Pullaiah & N. Yasoda	of Sps	R.R.V.Raju & T. Pullaiah	Sps			
1	Leguminosae	108	Leguminosae	97	Leguminosae	103	Leguminosae	156			
2	Poaceae	68	Euphorbiaceae	51	Graminae (Poaceae)	82	Poaceae	111			
3	Euphorbiaceae	52	Poaceae	48	Euphorbiaceae	35	Euphorbiaceae	54			
4	Acanthaceae	39	Acanthaceae	38	Asteraceae	35	Asteraceae	49			
5	Asteraceae	31	Asteraceae	33	Cyperaceae	33	Acanthaceae	45			
6	Rubiaceae	27	Rubiaceae	33	Acanthaceae	24	Cyperaceae	32			
7	Cyperaceae	22	Cyperaceae	27	Malvaceae	20	Convolvulaceae	30			
8	Asclepiadaceae	21	Malvaceae	27	Rubiaceae	19	Lamiaceae	27			
9	Lamiaceae	21	Convolvulaceae	23	Convolvulaceae	19	Rubiaceae	27			
10	Amaranthaceae	19	Amaranthaceae	17	Amaranthaceae	18	Malvaceae	22			

Table: 7. Comparision of Top 10 Genera with other district Floras of Andhra Pradesh

S.No.	Zoo Park	Sps	Ardhagiri	Sps	Chittoor (1991)	Sps	Chittoor (2008)	Sps
1	Euphorbia	15	Ficus	8	Crotalaria	18	Crotolaria	25
2	Justicia	12	Cassia	7	Cyperus	15	Indigofera	20
3	Cyperus	10	Cyperus	5	Ficus	13	Cyperus	19
4	Acacia	9	Ipomoea	5	Indigofera	13	Cassia	18
5	Crotalaria	9	Indigofera	5	Ipomoea	11	Euphorbia	17
6	Senna	9	Sida	5	Euphorbia	10	Ficus	17
7	Fimbristylis	9	Euphorbia	4	Fimbristylis	10	Ipomoea	16
8	Indigofera	8	Cissus	4	Phyllanthus	10	Acacia	15
9	Eragrostis	7			Rhynchosia	10	Jasminum	13
10	Ipomoea	7			Desmodium	8	Solanum	13
S.No.	Flora of A.P.	Sps	Anantapur (1989)	Sps	Kurnool (1995)	Sps	Nellore (2002)	Sps
1	Crotalaria	44	Euphorbia	14	Euphorbia	16	Indigofera	17
2	Cyperus	33	Cyperus	14	Ipomoea	15	Crotalaria	15
3	Euphorbia	30	Indigofera	12	Crotalaria	14	Cyperus	14
4	Ipomoea	24	Crotalaria	10	Indigofera	14	Cassia	11
5	Indigofera	24	Indigofera	10	Cyperus	11	Euphorbia	10
6	Fimbristylis	23	Fimbristylis	10	Phyllanthus	11	Hedyotis	10
7	Cassia	22	Cassia	9	Cassia	10	Ipomoea	10
8	Ficus	22	Acacia	9	Eragrostis	10	Phyllanthus	9
9	Leucas	20	Grewia	9	Grewia	10	Ficus	8
10	Grewia	18	Ipomoea	9	Acacia	9	Grewia	8

Graph: 3: Top 20 Genera of S.V. Zoological park




Table: 8. Monocots Vs Dicots % and Ratio of the Zoo Park with other regional floras

r]	ngi osperms	Т	D	%	M	%	R	T	D	%	M	%	R	T	D	%	M	%	R	
			No of Species						No of Genera						No of Families					
	A.P	2601	1927	74.09	674	25.91	2.86	1035	784	75.75	251	24.25	3.12	173	139	80.35	34	19.65	4.09	
	Zoo Park	702	639	81.71	143	18.29	4.47	482	398	82.57	84	17.43	4.74	108	88	18.48	20	18.51	4.40	
	Ardhagiri	299	245	81.94	54	18.06	4.54	210	177	84.29	33	15.71	5.36	59	54	91.53	5	8.47	10.80	
	Chittoor	983	813	82.71	168	17.09	4.84	552	457	82.79	95	17.21	4.81	136	112	82.35	24	17.65	4.67	
	Chittoor	1756	1391	79.21	365	20.79	3.81	879	694	78.95	185	21.05	3.75	176	137	77.84	39	22.16	3.51	
	Anantapur	707	554	78.36	153	21.64	3.62	398	320	80.40	78	19.60	4.10	101	84	83.17	17	16.83	4.94	
	Kurnool	1047	836	79.85	211	20.15	3.96	562	450	80.07	23	4.09	19.57	124	101	81.45	23	18.55	4.39	
	Nellore	926	792	85.53	134	14.47	5.91	525	429	81.71	96	18.29	4.47	142	119	83.80	23	16.20	5.17	

D: Dicots; M: Monocots; R: Ratio; T: Total

#### **DISCUSSION:**

Floristic enumeration of S.V.Zoological Park has been compared with the floristic studies of Andhra Pradesh and other significant floras of Rayalaseema (Chittoor, Kadapa, Kurnool and Nellore) Districts along with the flora of Ardhagiri hills of Chittoor District; Total number of dicot and monocot species, genera and families. (Table: 8). Revealed that the Andhra Pradesh flora resulted a total of 2601 species to that of 1927 (74.89%) dicotyledons and 674 (25.91%) monocotyledon species. Chittoor district represents a total of 1756 species with 1391 (79.21%) dicots and 365 (20.21%) monocotyledon species; followed by Kurnool district 1047 species with 836 (79.85 %) dicots 211 (20.15 %) monocots; Nellore district represents 926 species with 792 (85.53 %) dicots, 134 (14.47 %) monocots; Anantapur district with lowest number, 707 species to that of 544 (78.36 %) dicots and 153 (21.64 %) monocotyledon species. But it is observed that the distribution of species in the selected area S.V.Zoological Park showed more diversity with 782 Angiosperm species, 639 belongs to dicotyledons with 81.71 % and 143 monocotyledons with 18.29 %. When compared with the ratios between species, genera and families between monocots Vs dicots among other floras it is observed A.P Flora represents monocots Vs dicot species 1:2.86; between genera 1:3.12; between families 1:4.09; Ardhagiri flora 1:5.4 species; 1: 5.36 genera, 1: 10.80 families; Chittoor district 1: 3.81 species, 1 :3.75 genera; 1:3.51 families; Kurnool with 1:396 species; 1:4.02 genera and 1:4.39 families; Nellore district with 1:5.91 species, 1:4.47 genera, 1:5.17 families, Anantapur district with 1:3.62 species, 1:4.02 genera and 1:4.94 families. When compare with S.V.Zoological Park Monocots Vs Dicots species ratio 1:4.47; genera. 1:4.74 and between families 1:4.40 ratios showed its rich diversity compared to other floras of the state, and also shows close relationship with Chittoor, Kurnool and Nellore district floras in the species, genera and families distribution and diversity.

Hence it reveals the distribution of species in all district floras of Rayalaseema and the Andhra Pradesh as a whole is mainly dependent on the geographical and ecological conditions of the region, but not on the extent of area which the species are distributed. Chittoor and Kurnool district are located along the Eastern Ghats represents with rich floral diversity, further it is followed by Nellore district and least in the species distribution in Ananthapur district because of its geographical, ecological and climatic conditions leads for drought conditions to that of desert atmosphere. Whereas Ardhagiri hills which covers most of the area with rocks also represents proportional diversity in monocots Vs dicots equal to that of other floras even though the extent of area is very less than the representative Chittoor District. It signifies in the ratio and percentage between the families of dicots 54 (91.53%) and monocots 5 (8.47%) with 1:10.80 ratio denotes the less number of monocotyledon families with more number of genera and species diversity which may be possible with the grass species. Zoo Park flora signifies in the ratio of species, genera and families with almost equal as 1:4.47; 1:4.74 and 1:4.40 compared to that of Chittoor distinct flora except in the total number of species may be due to the small extent of area. But the diversity of the species is very rich with in the prevailed area in terms of Dicotyledons to monocotyledons. The aim of the floristic studies of a protected area resulted in a good number of species with dry deciduous spiny scrub vegetation along with patches of ever green shrubs and small tree species includes rare and endangered species of highly threatened tree Pterocarpus santalinus (Red sanders) in its natural habitat.

Many other species shown interest in the diversity as Capparis to be studied critically in particular to this region. Studies also necessiciated on the forage and food sources of natural birds and other native mammalian species to be conserved more in particular. Many interesting medicinal herbs are distributed in this locality as Hemidesmus, Tylophora, Decalepis, Gymnema, Holostemma, Ceropegia, Sarcostemma and Caralluma belongs to the family Asclepiadaceae as major species to be conserved for future generations. At the outset floristic enumeration of the S.V.Zoological Park revealed its significant distribution of Angiosperm species which are almost equal to that of the Chittoor district flora with in a small extent of area with much diversity of species. The reason may be the S.V.Zoological park is under conservation since 30 years, and also this area is a forest peripheral of S.V. National Park.

#### **CONCLUSION:**

It is recommended for the studies in identification of flora which are the major food source of native birds in different seasons; the species which attract butterflys; the major food source for various herbivorous species to be conserve and measures for the increase in the area of distribution of all above species.

#### **REFERENCES:**

- 1.IUCN List of National Parks and Protected Areas, IUCN Gland, United Nations, 1994.
- 2.IUCN. Parks for Life: Report of the IVth World Congress on National Parks and Protected Areas. Gland, Switzerland: IUCN. 1993
- 3.CNPPA, Parks for life; Action for protected areas in Europe, Workshop III.8, IUCN, Gland Switzerland., 1994.
- 4.http:/forest.ap.in/svzp-Introduction.htm.
- 5. Jain. S.K., Rao. R.R., A Handbook of field and Herbarium Today and Tomorrow printers and publishers, New Delhi, 1977.
- 6.Gamble, J.S. And C.E.S. Fischer. Flora of the Presidency of Madras. 1957. Vol.1-3, BSI Publication.
- 7. Matthew. K.M. Flora of the Tamilnadu, Carnatic 1-5 Volumes, Rapinat, Herbarium, St. Joseph's College, Tiruchirapalli.
- 8.Ellis. J.L. Contribution to the vascular flora of Bala palli range of Seshachalam Forest, Kadapa District, Andhra Pradesh. Bull. Bot. Surv. India. 1966, 8 (3&4); 435-437.
- 9. Venkata Raju, R.R. and Pullaiah, T. Flora of Kurnool District, Andhra Pradesh. Gajendra Singh Gahlot at Shiva Offset Press, and Doon Phototype Printers, for Bishen Singh Mahendra Pal Singh, Dehra Dun, India. 1995.
- 10. Pullaiah, T. and Yesoda, N. Flora of Anantapur District. Andhra Pradesh, India. Gajendra Singh Gahlot at Shiva Offset Press for Bishen Singh Mahendra Pal Singh Publishers, Dehra Dun, India. 1989.
- 11.Ranga Charyulu, D. Floristic Studies of Chittoor District of Andhra Pradesh, Ph.D. Thesis, Sri Venkateswara University, Tirupati. 1991.
- 12.Madhava Chetty, K., Sivaji, K. and Tulasi Rao, K. Flowering Plants of Chittoor District, Andhra Pradesh, India. Student Offset Printers, Tirupati, 2008.
- 13. Suryanarayana, B. and Sreenivasa Rao, A. Flora of Nellore District, Andhra Pradesh. Gurudev Prakashan Publishers, Wadala Mahadeo, Shrirampur Dist. Ahmednagar, Maharashtra, 2002.



## **A. Job Roger Binny**Dept of Botany, Sri Venkateswara University, Tirupati .

## Publish Research Article International Level Multidisciplinary Research Journal For All Subjects

Dear Sir/Mam,

We invite unpublished Research Paper, Summary of Research Project, Theses, Books and Book Review for publication, you will be pleased to know that our journals are

## Associated and Indexed, India

- ★ International Scientific Journal Consortium
- \* OPEN J-GATE

## Associated and Indexed, USA

- Google Scholar
- EBSCO
- DOAJ
- Index Copernicus
- Publication Index
- · Academic Journal Database
- Contemporary Research Index
- Academic Paper Databse
- Digital Journals Database
- Current Index to Scholarly Journals
- Elite Scientific Journal Archive
- Directory Of Academic Resources
- Scholar Journal Index
- Recent Science Index
- Scientific Resources Database
- Directory Of Research Journal Indexing

Indian Streams Research Journal 258/34 Raviwar Peth Solapur-413005,Maharashtra Contact-9595359435 E-Mail-ayisrj@yahoo.in/ayisrj2011@gmail.com Website: www.isrj.net